

Wetenschappelijke publicaties artrose en prothesiologie: knie

- Kornuijt A, van der Weegen W, van Drumpt R. Bewegen, sporten en werken met een totale knieprothese. Deel 3: Stijfheid van de knie na de operatie. *SPORTgericht* 2025;79(1):16-19.
- Theeuwen DMJ, Bemelmans YFL, Boonen B, Haveman I, van der Weegen W, Schotanus MGM. The influence of the number of postoperative radiological outliers on the survival and clinical outcome of total knee arthroplasty. *Journal of Clinical Orthopaedics and Trauma* 2025;60:102834.
- Theeuwen DMJ, Dorling IM, Most J, van Drumpt RAM, van der Weegen W, Welting TJM, Schotanus MGM, Boonen B. Patient-specific instrumentation improved clinical outcome and implant survival but is not superior compared to conventional total knee arthroplasty: ten years follow-up of a multicenter double-blind randomized controlled trial. *Knee Surg Sports Traumatol Arthrosc* 2025;33(4):1371-1377.
- Kornuijt A, van der Weegen W, van Drumpt R. Bewegen, sporten en werken met een totale knieprothese. Deel 1: Van operatie tot revalidatie. *SPORTgericht* 2024;78(4):44-48.
- Kornuijt A, van der Weegen W, van Zaanen Y, van Drumpt R. Bewegen, sporten en werken met een totale knieprothese. Deel 2: Terugkeer naar ADL, sport en werk. *SPORTgericht* 2024;78(6):28-33.
- Kornuijt A, van der Weegen W, Agricola R, Lenssen T. Satisfactory adherence to the Dutch physical therapy guideline for patients with a total knee arthroplasty. A survey study involving 103 specialised physical therapists. *Musculoskeletal Care* 2024;22(2):e1909.
- Kornuijt A, Bongers B, Marcellis R, Lenssen AF. Submaximal cardiopulmonary exercise testing to assess preoperative aerobic capacity in patients with knee osteoarthritis scheduled for total knee arthroplasty: a feasibility study. *Physiother Theory Pract* 2024;40(3):603-616.
- Kornuijt A, Kuijjer P, van Drumpt R, Siebelt M, Lenssen T, van der Weegen W. Hoge mate van fysieke activiteit na een totale knieprothese (TKP) lijkt geen nadeel. *FysioPraxis* 2023;32(1):25.
- van Schie P, van Bodegom-Vos L, Zijdemans TM, Nelissen RGHH, Marang-van de Mheen PJ, IQ Joint study group. Effectiveness of a multifaceted quality improvement intervention to improve patient outcomes after total hip and knee arthroplasty: a registry nested cluster randomised controlled trial. *BMJ Qual Saf* 2023;32:34-46.
- van der Weegen W, Das D, Vrints K, Agricola R, Siebelt M. A 20 mg dose of dexamethasone does not reduce the proportion of joint replacement patients needing rescue analgesia: a matched cohort study. *Ann Joint* 2023;8:4.

- van Eck J, Liu WY, Goosen JHM, Rijnen WHC, van der Zwaard BC, Heesterbeek P, van der Weegen W, The Further Members of Regional Prosthetic Joint Infection Group. Higher 1-year risk of implant removal for culture-positive than for culture-negative DAIR patients following 383 primary hip or knee arthroplasties. *J Bone Jt Infect* 2022;7(4):143-149.
- Kamp MC, Liu WY, Goosen JHM, Rijnen WHC, van Steenberg LN, van der Weegen W, Regional Prosthetic Joint Infection Working Group. Mismatch in capture of periprosthetic joint infections between the Dutch arthroplasty register (LROI) and a detailed regional periprosthetic joint infection registry. *J Arthroplasty* 2022;37(1):126-131.
- Kornuijt A, Kuijjer PPFM, van Drumpt RA, Siebelt M, Lenssen AF, van der Weegen W. A high physical activity level after total knee arthroplasty does not increase the risk of revision surgery during the first twelve years. A systematic review with meta-analysis and GRADE. *Knee* 2022;39:168-184.
- Wang Q, Runhaar J, Kloppenburg M, Boers M, Bijlsma JWJ, Bacardit J, Bierma-Zeinstra SMA, CREDO experts group. A machine learning approach reveals features related to clinicians' diagnosis of clinically relevant knee osteoarthritis. *Rheumatology* 2023;62(8):2732-2739.
- van der Weegen W, Sijbesma T, Siebelt M, Poolman RW. The effect of postponing joint replacement surgery due to COVID-19 on quality of life for patients suffering from end-stage osteoarthritis. *Wetenschap in Beweging* 2021;28:1.
- Kornuijt A, de Kort GJL, Das D, Lenssen AF, van der Weegen W. Recovery of knee range of motion after total knee arthroplasty in the first postoperative weeks: poor recovery can be detected early. *Musculoskelet Surg* 2019;103(3):289-297.
- Kornuijt A, van der Weegen W, de Kort J, Lenssen AF. De stijve totale knieprothese: snel herkennen en handelen! *FysioPraxis* 2019;28(9):24-27.
- Schotanus MGM, Boonen B, van Weegen W, Hoekstra H, van Drumpt R, Borghans R, Vos R, van Rhijn L, Kort NP. No difference in mid-term survival and clinical outcome between patient-specific and conventional instrumented total knee arthroplasty: a randomized controlled trial. *Knee Surg Sports Traumatol Arthrosc* 2019;27(5):1463-1468.
- Timmers T, Janssen L, van der Weegen W, Das D, Marijnissen WJ, Hannink G, van der Zwaard BC, Plat A, Thomassen B, Swen JW, Kool RB, Lambers Heerspink FO. The effect of an app for day-to-day postoperative care education on patients with total knee replacement: randomized controlled trial. *JMIR Mhealth Uhealth* 2019;7(10):e15323.
- van Zaanen Y, van Geenen RCI, Pahlplatz TMJ, Kievit AJ, Hoozemans MJM, Bakker EWP, Blankevoort L, Schafröth MU, Haverkamp D, Vervest TMJS, Das DHPW, van der Weegen W, Scholtes VA, Frings-Dresen MHW, Kuijjer PPFM. Three out of ten working patients expect no clinical improvement of their ability to perform work-related knee-demanding activities after total knee arthroplasty: a multicenter study. *J Occup Rehabil* 2019;29(3):585-594.
- Kornuijt A, Das D, Sijbesma T, de Vries L, van der Weegen W. Manipulation under anesthesia following total knee arthroplasty: a comprehensive review of literature. *Musculoskelet Surg* 2018;102(3):223-230.

- Timmers T, Janssen L, Pronk Y, van der Zwaard BC, Koëter S, van Oostveen D, de Boer S, Kremers K, Rutten S, Das D, van Geenen RC, Koenraadt KL, Kusters R, van der Weegen W. Assessing the efficacy of an educational smartphone or tablet app with subdivided and interactive content to increase patients' medical knowledge: randomized controlled trial. *JMIR Mhealth Uhealth* 2018;6(12):e10742.
- van der Weegen W, Verduin CM, Graumans M, Hoekstra HJ. Successful debridement of a knee joint prosthesis infected with *Listeria Monocytogenes*. Case report and review of current literature. *J Bone Jt Infect* 2018;3(4):203-206.
- Bastick AN, Damen J, Agricola R, Brouwer RW, Bindels PJ, Bierma-Zeinstra SM. Characteristics associated with joint replacement in early symptomatic knee or hip osteoarthritis: 6-year results from a nationwide prospective cohort study (CHECK). *Br J Gen Pract* 2017;67(663):e724-e731.
- Löwik C, Wagenaar FC, Stevens M, Bulstra S, Pronk Y, van den Akker-Scheek I, Wouthuyzen-Bakker M, Nelissen R, Poolman RW, van der Weegen W, Jutte PC. Managing persistent wound leakage after total knee and hip arthroplasty. Results of a nationwide survey among Dutch orthopedic surgeons. *J Bone Jt Infect*. 2017;(4):202-207.
- Löwik CAM, Wagenaar FC, van der Weegen W, Poolman RW, Nelissen RGHH, Bulstra SK, Pronk Y, Vermeulen KM, Wouthuyzen-Bakker M, van den Akker-Scheek I, Stevens M, Jutte PC, LEAK study group. LEAK study: design of a nationwide randomised controlled trial to find the best way to treat wound leakage after primary hip and knee arthroplasty. *BMJ Open* 2017;7(12):e018673.
- Boonen B, Schotanus MGM, Kerens B, van der Weegen W, Hoekstra HJ, Kort NP. No difference in clinical outcome between patient-matched positioning guides and conventional instrumented total knee arthroplasty two years post-operatively: a multicentre, double-blind, randomised controlled trial. *Bone Joint J* 2016;98-B(7):939-944.
- van Drumpt RAM, van der Weegen W, King W, Toler K, Macenski MM. Safety and treatment effectiveness of a single autologous protein solution injection in patients with knee osteoarthritis. *Biores Open Access* 2016;5(1):261-268.
- King W, van der Weegen W, van Drumpt RAM, Scoons H, Toler K, Woodell-May J. White blood cell concentration correlates with increased concentrations of IL-1ra and improvement in WOMAC pain scores in an open-label safety study of autologous protein solution. *J Exp Orthop* 2016;3(1):9.
- van Tiel J, Siebelt M, Reijman M, Bos PK, Waarsing JH, Zuurmond AM, Nasserinejad K, van Osch GJ, Verhaar JA, Krestin GP, Weinans H, Oei EH. Quantitative in vivo CT arthrography of the human osteoarthritic knee to estimate cartilage sulphated glycosaminoglycan content: correlation with ex-vivo reference standards. *Osteoarthritis Cartilage* 2016;24(6):1012-1020.
- de Vries LMA, van der Weegen W, Neve WC, Das HPW, Ridwan BU, Steens J. The effectiveness of debridement, antibiotics and irrigation for periprosthetic joint infections after hip or knee arthroplasty. A 15 years retrospective study in two community hospitals in the Netherlands. *J Bone Joint Infect* 2016;1:20-24.
- Glyn-Jones S, Palmer AJ, Agricola R, Price AJ, Vincent TL, Weinans H, Carr AJ. Osteoarthritis. *Lancet* 2015;386(9991):376-387.

- Siebelt M, Korthagen N, Wei W, Groen H, Bastiaansen-Jenniskens Y, Müller C, Waarsing JH, de Jong M, Weinans H. Triamcinolone acetonide activates an anti-inflammatory and folate receptor-positive macrophage that prevents osteophytosis in vivo. *Arthritis Res Ther* 2015;17:352.
- van Spil WE, Agricola R, Drossaers-Bakker KW, Weinans H, Lafeber FP. Associations of markers of matrix metabolism, inflammation markers, and adipokines with superior cam deformity of the hip and their relation with future hip osteoarthritis. *Osteoarthritis Cartilage* 2015;23(11):1897-1905.
- van der Weegen W, Wullems JA, Bos E, Noten H, van Drumpt RAM. No difference between intra-articular injection of hyaluronic acid and placebo for mild to moderate knee osteoarthritis: a randomized, controlled, double-blind trial. *J Arthroplasty* 2015;30(5):754-757.
- van Buul GM, Siebelt M, Leijns MJ, Bos PK, Waarsing JH, Kops N, Weinans H, Verhaar JA, Bernsen MR, van Osch GJ. Mesenchymal stem cells reduce pain but not degenerative changes in a mono-iodoacetate rat model of osteoarthritis. *J Orthop Res* 2014;32(9):1167-1174.
- Siebelt M, Groen HC, Koelewijn SJ, de Blois E, Sandker M, Waarsing JH, Müller C, van Osch GJ, de Jong M, Weinans H. Increased physical activity severely induces osteoarthritic changes in knee joints with papain induced sulfate-glycosaminoglycan depleted cartilage. *Arthritis Res Ther* 2014;16(1):R32.
- Siebelt M, Waarsing JH, Groen HC, Müller C, Koelewijn SJ, de Blois E, Verhaar JA, de Jong M, Weinans H. Inhibited osteoclastic bone resorption through alendronate treatment in rats reduces severe osteoarthritis progression. *Bone* 2014;66:163-170.
- Siebelt M, van der Windt AE, Groen HC, Sandker M, Waarsing JH, Müller C, de Jong M, Jahr H, Weinans H. FK506 protects against articular cartilage collagenous extra-cellular matrix degradation. *Osteoarthritis Cartilage* 2014;22(4):591-600.
- Austen S, van der Weegen W, Verduin CM, van der Valk M, Hoekstra HJ. Coccidioidomycosis infection of a total knee arthroplasty in a nonendemic region. *J Arthroplasty* 2013;28(2):375.e13-15.
- Boonen B, Schotanus MG, Kerens B, van der Weegen W, van Drumpt RA, Kort NP. Intra-operative results and radiological outcome of conventional and patient-specific surgery in total knee arthroplasty: a multicentre, randomised controlled trial. *Knee Surg Sports Traumatol Arthrosc* 2013;21(10):2206-2212.
- Koolen NEH, van Drumpt R, Hoekstra HJ, van der Weegen W. Recovery after unicompartmental knee arthroplasty. An observational pilot study on the need for post-operative physiotherapy. *Nederlands Tijdschrift voor Orthopaedie* 2013;20(4):101-105.
- Siebelt M, Jahr H, Groen HC, Sandker M, Waarsing JH, Kops N, Müller C, van Eden W, de Jong M, Weinans H. Hsp90 inhibition protects against biomechanically induced osteoarthritis in rats. *Arthritis Rheum* 2013;65(8):2102-2112.

- van Tiel J, Siebelt M, Waarsing JH, Piscaer TM, van Straten M, Booij R, Dijkshoorn ML, Kleinrensink GJ, Verhaar JA, Krestin GP, Weinans H, Oei EH. CT arthrography of the human knee to measure cartilage quality with low radiation dose. *Osteoarthritis Cartilage* 2012;20(7):678-685.
- Weinans H, Siebelt M, Agricola R, Botter SM, Piscaer TM, Waarsing JH. Pathophysiology of peri-articular bone changes in osteoarthritis. *Bone* 2012;51(2):190-196.
- Miclea RL, Siebelt M, Finos L, Goeman JJ, Löwik CW, Oostdijk W, Weinans H, Wit JM, Robanus-Maandag EC, Karperien M. Inhibition of Gsk3 β in cartilage induces osteoarthritic features through activation of the canonical Wnt signaling pathway. *Osteoarthritis Cartilage* 2011;19(11):1363-1372.
- Siebelt M, van Tiel J, Waarsing JH, Piscaer TM, van Straten M, Booij R, Dijkshoorn ML, Kleinrensink GJ, Verhaar JA, Krestin GP, Weinans H, Oei EH. Clinically applied CT arthrography to measure the sulphated glycosaminoglycan content of cartilage. *Osteoarthritis Cartilage* 2011;19(10):1183-1189.
- Siebelt M, Waarsing JH, Kops N, Piscaer TM, Verhaar JA, Oei EH, Weinans H. Quantifying osteoarthritic cartilage changes accurately using in vivo microCT arthrography in three etiologically distinct rat models. *J Orthop Res* 2011;29(11):1788-1794.
- Broekhuis D, van der Weegen W, de Sain K, Kort NP. Röntgenfoto's bij unicompartimentele knieprotheses, het voordeel van positionering onder doorlichting. *Nederlands Tijdschrift voor Orthopaedie* 2010;17(1):31-36.