

Wetenschappelijke publicaties artrose en prothesiologie: heup

- Boel F, Wortel J, van Buuren MMA, Rivadeneira F, van Meurs JBJ, Runhaar J, Bierma-Zeinstra SMA, Agricola R. DXA images vs. pelvic radiographs: reliability of hip morphology measurements. *Osteoarthritis Cartilage* 2025;33(2):283-292.
- Gijon-Nogureon G, Balint P, Batalov A, Ostojic P, Sollman N, van Middelkoop M, Agricola R, Naili JE, Milovanovic D, Popova S, Kazakova M, Nuernberger S, Aulin C, Karalilova R, Henrotin Y. Terminologies and definitions used to classify patients with osteoarthritis: a scoping review. *BMC Rheumatol* 2025;9(1):32.
- Riedstra NS, Boel F, van Buuren MMA, Ahedi H, Arbabi V, Arden N, Baart SJ, Bierma-Zeinstra SMA, Cicuttini FM, Cootes TF, Crossley KM, Felson DT, Giellis WP, Heerey J, Jones G, Kluzek S, Lane LE, Lindner C, Lynch JA, van Meurs JBJ, Mosler A, Nelson AE, Nevitt MC, Oei EH, Runhaar J, Tang J, Weinans H, Agricola R. Acetabular dysplasia and the risk of developing hip osteoarthritis within 4-8 years: an individual participant data meta-analysis of 18,807 hips from the World COACH consortium. *Osteoarthritis Cartilage* 2025;33(3):373-382.
- Agricola R, van Buuren MMA, Kemp JL, Weinans H, Runhaar J, Bierma-Zeinstra SMA. Femoroacetabular impingement syndrome in middle-aged individuals is strongly associated with the development of hip osteoarthritis within 10-year follow-up: a prospective cohort study (CHECK). *Br J Sports Med* 2024;58(18):1061-1067.
- van Buuren MMA, Riedstra NS, van den Berg MA, Boel FDEM, Ahedi H, Arbabi V, Arden NK, Bierma-Zeinstra SMA, Boer CG, Cicuttini F, Cootes TF, Crossley K, Felson D, Gielis WP, Heerey J, Jones G, Kluzek S, Lane NE, Lindner C, Lynch JA, van Meurs J, Mosler AB, Nelson AE, Nevitt M, Oei E, Runhaar J, Tang J, Weinans H, Agricola R. Cohort profile: Worldwide Collaboration on OsteoArthritis prediction for the Hip (World COACH) - an international consortium of prospective cohort studies with individual participant data on hip osteoarthritis. *BMJ Open* 2024;14(4):e077907.
- Heerey JJ, van Klij P, Agricola R, Dijkstra HP, Plass L, Crossley KM, Kemp JL. Preventing hip osteoarthritis in athletes: is it really a mission impossible? *Br J Sports Med* 2024;58(9):465-467.
- Riedstra NS, Boel F, van Buuren M, Eygendaal D, Bierma-Zeinstra S, Runhaar J, Agricola R. Pincer morphology is not associated with hip osteoarthritis unless hip pain is present. *Arthritis Care Res (Hoboken)* 2024;76(5):644-651.
- Tang J, Boel F, van Buuren MMA, Riedstra NS, Runhaar J, Bierma-Zeinstra S, Agricola R. The different subtypes of cam morphology as defined by statistical shape modeling and their relationship with the development of hip osteoarthritis: a nationwide prospective cohort study (CHECK) with 10 years follow-up. *Osteoarthritis Cartilage* 2024;32(12):1647-1654.
- van der Weegen W, Warren T, Das D, Agricola R, Timmers T, Siebelt M. Operative or nonoperative treatment is predicted accurately for patients who have hip complaints consulting an orthopaedic surgeon using machine learning algorithms trained with prehospital acquired history-taking data. *J Arthroplasty* 2024;39(5):1173-1177.

- Kim JS, Agricola R, Kim YJ, Lane NE, Millis MB, Nelson AE, Runhaar J, Shefelbine SJ, Bostrom MP. Arthritis Foundation/HSS Workshop on hip osteoarthritis, part 1: epidemiology, early development, and cohorts from around the world. *HSS J* 2023;19(4):395-401.
- Pronk Y, van der Weegen W, Schreurs BW, Pilot P. Quality of total hip arthroplasty health care based on four years of patient-reported outcomes in the Netherlands. *Health Qual Life Outcomes* 2023;21(1):24.
- Riedstra NS, Vinge R, Herfkens J, Eygendaal D, Bierma-Zeinstra SMA, Runhaar J, van Buuren MMA, Agricola R. Acetabular dysplasia and the risk of developing hip osteoarthritis at 2, 5, 8, and 10 years follow-up in a prospective nationwide cohort study (CHECK). *Semin Arthritis Rheum* 2023;60:152194.
- Runhaar J, van Berkel AC, Agricola R, van Meurs J, Bierma-Zeinstra SMA. Risk factors and population-attributable fractions for incident hip osteoarthritis. *HSS J* 2023;19(4):407-412.
- van Schie P, van Bodegom-Vos L, Zijdemans TM, Nelissen RGHH, Marang-van de Mheen PJ, IQ Joint study group. Effectiveness of a multifaceted quality improvement intervention to improve patient outcomes after total hip and knee arthroplasty: a registry nested cluster randomised controlled trial. *BMJ Qual Saf* 2023;32:34-46.
- Vinge R, Riedstra N, Tiderius CJ, Bierma-Zeinstra S, Agricola R, Runhaar J. Hip dysplasia as risk factor for clinically relevant and radiographic hip osteoarthritis: 10-year results from the CHECK cohort. *Rheumatology (Oxford)* 2023, online ahead of print.
- van der Weegen W, Das D, Vrints K, Agricola R, Siebelt M. A 20 mg dose of dexamethasone does not reduce the proportion of joint replacement patients needing rescue analgesia: a matched cohort study. *Ann Joint* 2023;8:4.
- Ahedi H, Winzenberg T, Bierma-Zeinstra S, Blizzard L, van Middelkoop M, Agricola R, Waarsing JH, Cicuttini F, Jones G. A prospective cohort study on cam morphology and its role in progression of osteoarthritis. *Int J Rheum Dis* 2022;25(5):601-612.
- van Eck J, Liu WY, Goosen JHM, Rijnen WHC, van der Zwaard BC, Heesterbeek P, van der Weegen W, The Further Members of Regional Prosthetic Joint Infection Group. Higher 1-year risk of implant removal for culture-positive than for culture-negative DAIR patients following 383 primary hip or knee arthroplasties. *J Bone Jt Infect* 2022;7(4):143-149.
- Kamp MC, Liu WY, Goosen JHM, Rijnen WHC, van Steenberghe LN, van der Weegen W, Regional Prosthetic Joint Infection Working Group. Mismatch in capture of periprosthetic joint infections between the Dutch arthroplasty register (LROI) and a detailed regional periprosthetic joint infection registry. *J Arthroplasty* 2022;37(1):126-131.
- van der Weegen W, Hoekstra H, Brakel K, Sijbesma T. Limited need for screening of metal-on-metal hip resurfacing patients beyond 10 years of follow-up. *Hip Int* 2022;32(1):106-112.
- van Buuren MMA, Arden NK, Bierma-Zeinstra SMA, Bramer WM, Casartelli NC, Felson DT, Jones G, Lane NE, Lindner C, Maffiuletti NA, van Meurs JBJ, Nelson AE, Nevitt MC, Valenzuela PL, Verhaar JAN, Weinans H, Agricola R. Statistical shape modeling of the hip and the association with hip osteoarthritis: a systematic review. *Osteoarthritis Cartilage* 2021;29(5):607-618.

- Casartelli NC, Maffiuletti NA, Valenzuela PL, Grassi A, Ferrari E, van Buuren MMA, Nevitt MC, Leunig M, Agricola R. Is hip morphology a risk factor for developing hip osteoarthritis? A systematic review with meta-analysis. *Osteoarthritis Cartilage* 2021;29(9):1252-1264.
- Heerey JJ, Srinivasan R, Agricola R, Smith A, Kemp JL, Pizzari T, King MG, Lawrenson PR, Scholes MJ, Souza RB, Link T, Majumdar S, Crossley KM. Prevalence of early hip OA features on MRI in high-impact athletes. The femoroacetabular impingement and hip osteoarthritis cohort (FORCe) study. *Osteoarthritis Cartilage* 2021;29(3):323-334.
- Kemp JL, Østerås N, Mathiessen A, Nordsletten L, Agricola R, Waarsing JH, Heerey JJ, Risberg MA. Relationship between cam morphology, hip symptoms, and hip osteoarthritis: the Musculoskeletal pain in Ullersaker Study (MUST) cohort. *Hip Int* 2021;31(6):789-796.
- Pronk Y, Pilot P, van der Weegen W, Brinkman JM, Schreurs BW. A patient-reported outcome tool to triage total hip arthroplasty patients to hospital or video consultation: pilot study with expert panels and a cohort of 1228 patients. *JMIR Form Res* 2021;5(12):e31232.
- Siebelt M, Das D, van den Moosdijk A, Warren T, van der Putten P, van der Weegen W. Machine learning algorithms trained with pre-hospital acquired history taking data can accurately differentiate diagnoses in patients with hip complaints. *Acta Orthop* 2021;92(3):254-257.
- Wang Q, Runhaar J, Kloppenburg M, Boers M, Bijlsma JWJ, Bierma-Zeinstra SMA; CREDO expert group. Diagnosis of early stage knee osteoarthritis based on early clinical course: data from the CHECK cohort. *Arthritis Res Ther* 2021;23(1):217.
- van der Weegen W, Sijbesma T, Siebelt M, Poolman RW. The effect of postponing joint replacement surgery due to COVID-19 on quality of life for patients suffering from end-stage osteoarthritis. *Wetenschap in Beweging* 2021;28:1.
- Beddows TPA, van Klij P, Agricola R, Tak IJR, Piscaer T, Verhaar JAN, Weir A. Normal values for hip muscle strength and range of motion in elite, sub-elite and amateur male field hockey players. *Phys Ther Sport* 2020;46:169-176.
- Gielis WP, Weinans H, Welsing PMJ, van Spil WE, Agricola R, Cootes TF, de Jong PA, Lindner C. An automated workflow based on hip shape improves personalized risk prediction for hip osteoarthritis in the CHECK study. *Osteoarthritis Cartilage* 2020;28(1):62-70.
- Pronk Y, van der Weegen W, Vos R, Brinkman JM, van Heerwaarden RJ, Pilot P. What is the minimum response rate on patient-reported outcome measures needed to adequately evaluate total hip arthroplasties? *Health Qual Life Outcomes* 2020;18(1):379.
- Wang Q, Runhaar J, Kloppenburg M, Boers M, Bijlsma JWJ, Bierma-Zeinstra SMA, The CREDO Expert Group. The added value of radiographs in diagnosing knee osteoarthritis is similar for general practitioners and secondary care physicians; data from the CHECK early osteoarthritis cohort. *J Clin Med* 2020;9(10):3374.
- Heerey JJ, Kemp JL, Mosler AB, Jones DM, Pizzari T, Scholes MJ, Agricola R, Crossley KM. What is the prevalence of hip intra-articular pathologies and osteoarthritis in active athletes with hip and groin pain compared with those without? A systematic review and meta-analysis. *Sports Med* 2019;49(6):951-972.

- Heerey JJ, Kemp JL, Mosler AB, Jones DM, Pizzari T, Scholes MJ, Agricola R, Crossley KM. Correction to: What is the prevalence of hip intra-articular pathologies and osteoarthritis in active athletes with hip and groin pain compared with those without? A systematic review and meta-analysis. *Sports Med* 2019;49(7):1139-1141.
- Hirvasniemi J, Gielis WP, Arbabi S, Agricola R, van Spil WE, Arbabi V, Weinans H. Bone texture analysis for prediction of incident radiographic hip osteoarthritis using machine learning: data from the Cohort Hip and Cohort Knee (CHECK) study. *Osteoarthritis Cartilage* 2019;27(6):906-914.
- van der Weegen W, Kornuijt A, Das D, Vos R, Sijbesma T. It is safe to use minimal restrictions following posterior approach total hip arthroplasty: results from a large cohort study. *Hip Int* 2019;29(6):572-577.
- Crossley KM, Pandey MG, Majumdar S, Smith AJ, Agricola R, Semciw AI, Kemp JL, Heerey JJ, King MG, Lawrenson PR, Lin YC, Souza RB, Mosler AB, Link TM, Srinivasan R, Schache AG. Femoroacetabular impingement and hip Osteoarthritis Cohort (FORCe): protocol for a prospective study. *J Physiother* 2018;64(1):55.
- Kemp JL, Crossley KM, Agricola R, Geuskens F, van Middelkoop M. Radiographic hip osteoarthritis is prevalent, and is related to cam deformity 12-24 months post-hip arthroscopy. *Int J Sports Phys Ther* 2018;13(2):177-184.
- Bastick AN, Damen J, Agricola R, Brouwer RW, Bindels PJ, Bierma-Zeinstra SM. Characteristics associated with joint replacement in early symptomatic knee or hip osteoarthritis: 6-year results from a nationwide prospective cohort study (CHECK). *Br J Gen Pract* 2017;67(663):e724-e731.
- Kornuijt A, Das D, Sijbesma T, van der Weegen W. Niet méér luxaties door minimale leefregels na THP. Standaard leefregels versus minimale leefregels bij posterieure benadering. *FysioPraxis* 2017;26(2):30-33.
- Löwik C, Wagenaar FC, Stevens M, Bulstra S, Pronk Y, van den Akker-Scheek I, Wouthuyzen-Bakker M, Nelissen R, Poolman RW, van der Weegen W, Jutte PC. Managing persistent wound leakage after total knee and hip arthroplasty. Results of a nationwide survey among Dutch orthopedic surgeons. *J Bone Jt Infect*. 2017;(4):202-207.
- Löwik CAM, Wagenaar FC, van der Weegen W, Poolman RW, Nelissen RGHH, Bulstra SK, Pronk Y, Vermeulen KM, Wouthuyzen-Bakker M, van den Akker-Scheek I, Stevens M, Jutte PC, LEAK study group. LEAK study: design of a nationwide randomised controlled trial to find the best way to treat wound leakage after primary hip and knee arthroplasty. *BMJ Open* 2017;7(12):e018673.
- Saberi Hosnijeh F, Zuiderwijk ME, Versteeg M, Smeele HT, Hofman A, Uitterlinden AG, Agricola R, Oei EH, Waarsing JH, Bierma-Zeinstra SM, van Meurs JB. Cam deformity and acetabular dysplasia as risk factors for hip osteoarthritis. *Arthritis Rheumatol* 2017;69(1):86-93.
- van der Weegen W, Kornuijt A, Das D. Zijn luxaties van een totale heupprothese te voorkomen door leefregels? Postoperatieve leefregelprotocollen onnodig beperkend voor de patiënt. *FysioPraxis* 2017;26(2):24-29.
- Das D, van der Weegen W, Wullems J, Brakel K, Nelissen R. Peri-prosthetic pathology in 'at-risk' ceramic-on-polyethylene total hip arthroplasty: a clinical study using MARS-MRI in 50 patients. *Hip Int* 2016;26(3):244-228.

- Kornuijt A, Das D, Sijbesma T, van der Weegen W. The rate of dislocation is not increased when minimal precautions are used after total hip arthroplasty using the posterolateral approach: a prospective, comparative safety study. *Bone Joint J* 2016;98-B(5):589-594.
- Roze RH, Bierma-Zeinstra SM, Agricola R, Oei EH, Waarsing JH. Differences in MRI features between two different osteoarthritis subpopulations: data from the Osteoarthritis Initiative. *Osteoarthritis Cartilage* 2016;24(5):822-826.
- de Vries LMA, van der Weegen W, Neve WC, Das HPW, Ridwan BU, Steens J. The effectiveness of debridement, antibiotics and irrigation for periprosthetic joint infections after hip or knee arthroplasty. A 15 years retrospective study in two community hospitals in the Netherlands. *J Bone Joint Infect* 2016;1:20-24.
- van der Weegen W, Kornuijt A, Das D. Do lifestyle restrictions and precautions prevent dislocation after total hip arthroplasty? A systematic review and meta-analysis of the literature. *Clin Rehabil* 2016;30(4):329-339.
- Agricola R, Leyland KM, Bierma-Zeinstra SM, Thomas GE, Emans PJ, Spector TD, Weinans H, Waarsing JH, Arden NK. Validation of statistical shape modelling to predict hip osteoarthritis in females: data from two prospective cohort studies (Cohort Hip and Cohort Knee and Chingford). *Rheumatology (Oxford)* 2015;54(11):2033-2041.
- Glyn-Jones S, Palmer AJ, Agricola R, Price AJ, Vincent TL, Weinans H, Carr AJ. Osteoarthritis. *Lancet* 2015;386(9991):376-387.
- Siebelt M, Korthagen N, Wei W, Groen H, Bastiaansen-Jenniskens Y, Müller C, Waarsing JH, de Jong M, Weinans H. Triamcinolone acetonide activates an anti-inflammatory and folate receptor-positive macrophage that prevents osteophytosis in vivo. *Arthritis Res Ther* 2015;17:352.
- van Spil WE, Agricola R, Drossaers-Bakker KW, Weinans H, Lafeber FP. Associations of markers of matrix metabolism, inflammation markers, and adipokines with superior cam deformity of the hip and their relation with future hip osteoarthritis. *Osteoarthritis Cartilage* 2015;23(11):1897-1905.
- van Buul GM, Siebelt M, Leijts MJ, Bos PK, Waarsing JH, Kops N, Weinans H, Verhaar JA, Bernsen MR, van Osch GJ. Mesenchymal stem cells reduce pain but not degenerative changes in a mono-iodoacetate rat model of osteoarthritis. *J Orthop Res* 2014;32(9):1167-1174.
- Siebelt M, Agricola R, Weinans H, Kim YJ. The role of imaging in early hip OA. *Osteoarthritis Cartilage* 2014;22(10):1470-1480.
- Siebelt M, Waarsing JH, Groen HC, Müller C, Koelewijn SJ, de Blois E, Verhaar JA, de Jong M, Weinans H. Inhibited osteoclastic bone resorption through alendronate treatment in rats reduces severe osteoarthritis progression. *Bone* 2014;66:163-170.
- Siebelt M, van der Windt AE, Groen HC, Sandker M, Waarsing JH, Müller C, de Jong M, Jahr H, Weinans H. FK506 protects against articular cartilage collagenous extra-cellular matrix degradation. *Osteoarthritis Cartilage* 2014;22(4):591-600.
- de Vries LAM, van der Weegen W, Pilot P, Stolarczyk PA, Sijbesma T, Hoffman EL. The predictive value of radiostereometric analysis for stem survival in total hip arthroplasty. A systematic review. *Hip Int* 2014;24(3):215-222.

- van der Weegen W, Brakel K, Horn RJ, Wullems JA, Das D, Pilot P, Nelissen RGHH. Comparison of three different pseudotumour grading systems within a single cohort of metal-on-metal hip arthroplasty patients. *Skeletal Radiol* 2014;43(2):149-155.
- van der Weegen W, Sijbesma T, Hoekstra HJ, Brakel K, Pilot P, Nelissen RGHH. Treatment of pseudotumors after metal-on-metal hip resurfacing based on magnetic resonance imaging, metal ion levels and symptoms. *J Arthroplasty* 2014;29(2):416-421.
- Agricola R, Heijboer MP, Bierma-Zeinstra SM, Verhaar JA, Weinans H, Waarsing JH. Cam impingement causes osteoarthritis of the hip: a nationwide prospective cohort study (CHECK). *Ann Rheum Dis* 2013;72(6):918-923.
- Agricola R, Heijboer MP, Roze RH, Reijman M, Bierma-Zeinstra SM, Verhaar JA, Weinans H, Waarsing JH. Pincer deformity does not lead to osteoarthritis of the hip whereas acetabular dysplasia does: acetabular coverage and development of osteoarthritis in a nationwide prospective cohort study (CHECK). *Osteoarthritis Cartilage* 2013;21(10):1514-1521.
- Agricola R, Reijman M, Bierma-Zeinstra SM, Verhaar JA, Weinans H, Waarsing JH. Total hip replacement but not clinical osteoarthritis can be predicted by the shape of the hip: a prospective cohort study (CHECK). *Osteoarthritis Cartilage* 2013;21(4):559-564.
- Agricola R, Waarsing JH, Arden NK, Carr AJ, Bierma-Zeinstra SM, Thomas GE, Weinans H, Glyn-Jones S. Cam impingement of the hip: a risk factor for hip osteoarthritis. *Nat Rev Rheumatol* 2013;9(10):630-634.
- Sankar WN, Nevitt M, Parvizi J, Felson DT, Agricola R, Leunig M. Femoroacetabular impingement: defining the condition and its role in the pathophysiology of osteoarthritis. *J Am Acad Orthop Surg* 2013;21 Suppl 1:S7-S15.
- Siebelt M, Jahr H, Groen HC, Sandker M, Waarsing JH, Kops N, Müller C, van Eden W, de Jong M, Weinans H. Hsp90 inhibition protects against biomechanically induced osteoarthritis in rats. *Arthritis Rheum* 2013;65(8):2102-2112.
- van der Weegen W, Brakel K, Horn RJ, Hoekstra HJ, Sijbesma T, Pilot P, Nelissen RGHH. Asymptomatic pseudotumours after metal-on-metal hip resurfacing show little change within one year. *Bone Joint J* 2013;95-B(12):1626-1231.
- van der Weegen W, Smolders JMH, Sijbesma T, Hoekstra HJ, Brakel K, van Susante JLC. High incidence of pseudotumors after hip resurfacing even in low risk patients; results from an intensified MRI screening protocol. *Hip Int* 2013;23(3):243-249.
- van der Weegen W, Austen S, Sijbesma T, Hoekstra HJ. Polyethyleen slijtage van ongecementeerde cups – een analyse van 200 Mallory Head heupprotheses. *Nederlands Tijdschrift voor Orthopaedie* 2012;19(1):4-9.
- van der Weegen W, Hoekstra HJ, Sijbesma T, Austen S, Bos E, Poolman RW. Hip resurfacing in a district general hospital: 6-year clinical results using the ReCap hip resurfacing system. *BMC Musculoskelet Disord* 2012;13:247.
- Weinans H, Siebelt M, Agricola R, Botter SM, Piscaer TM, Waarsing JH. Pathophysiology of peri-articular bone changes in osteoarthritis. *Bone* 2012;51(2):190-196.

- Miclea RL, Siebelt M, Finos L, Goeman JJ, Löwik CW, Oostdijk W, Weinans H, Wit JM, Robanus-Maandag EC, Karperien M. Inhibition of Gsk3 β in cartilage induces osteoarthritic features through activation of the canonical Wnt signaling pathway. *Osteoarthritis Cartilage* 2011;19(11):1363-1372.
- Siebelt M, Waarsing JH, Kops N, Piscaer TM, Verhaar JA, Oei EH, Weinans H. Quantifying osteoarthritic cartilage changes accurately using in vivo microCT arthrography in three etiologically distinct rat models. *J Orthop Res* 2011;29(11):1788-1794.
- Siebelt M, van Tiel J, Waarsing JH, Piscaer TM, van Straten M, Booij R, Dijkshoorn ML, Kleinrensink GJ, Verhaar JA, Krestin GP, Weinans H, Oei EH. Clinically applied CT arthrography to measure the sulphated glycosaminoglycan content of cartilage. *Osteoarthritis Cartilage* 2011;19(10):1183-1189.
- Sybesma T, Röckmann H, van der Weegen W. Severe anaphylactic reaction to chlorhexidine during total hip replacement surgery. A case report. *Hip Int* 2011;21(5):630-632.
- van der Weegen W, Hoekstra HJ, Sijbesma T, Bos E, Schemitsch EH, Poolman RW. Survival of metal-on-metal hip resurfacing arthroplasty: a systematic review of the literature. *J Bone Joint Surg Br* 2011;93(3):298-306.